

Spikes in acute:chronic workload ratio (ACWR) associated with a 5-7 times greater injury rate in English Premier League football players

Bowen, Laura; Gross, Aleksander Stephan; Gimpel, Mo; Bruce-Low, Stewart; Li, Francois Xavier

DOI:

[10.1136/bjsports-2018-099422](https://doi.org/10.1136/bjsports-2018-099422)

License:

Creative Commons: Attribution-NonCommercial (CC BY-NC)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Bowen L, Gross AS, Gimpel M, et al Spikes in acute:chronic workload ratio (ACWR) associated with a 5–7 times greater injury rate in English Premier League football players: A comprehensive 3-year study', *British Journal of Sports Medicine*. <https://doi.org/10.1136/bjsports-2018-099422>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 26/06/2019

Bowen L, Gross AS, Gimpel M, et al Spikes in acute:chronic workload ratio (ACWR) associated with a 5–7 times greater injury rate in English Premier League football players: a comprehensive 3-year study *British Journal of Sports Medicine* Published Online First: 21 February 2019. doi: 10.1136/bjsports-2018-099422

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



OPEN ACCESS

Spikes in acute:chronic workload ratio (ACWR) associated with a 5–7 times greater injury rate in English Premier League football players: a comprehensive 3-year study

Laura Bowen,^{1,2} Aleksander Stephan Gross,² Mo Gimpel,³ Stewart Bruce-Low,³ Francois-Xavier Li¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2018-099422>).

¹Sport, Exercise and Rehabilitation Sciences, University of Birmingham, Birmingham, UK

²Southampton Football Club, Sports Medicine and Science, Southampton, UK

³Performance Support, Southampton Football Club, Southampton, UK

Correspondence to

Miss. Laura Bowen, Sport, Exercise and Rehabilitation Sciences, University of Birmingham, Birmingham B15 2TT, UK; lbowen@saintsfc.co.uk

Accepted 23 January 2019

ABSTRACT

Objectives We examined the relation between global positioning system (GPS)-derived workloads and injury in English Premier League football players (n=33) over three seasons.

Methods Workload and injury data were collected over three consecutive seasons. Cumulative (1-weekly, 2-weekly, 3-weekly and 4-weekly) loads in addition to acute:chronic workload ratios (ACWR) (acute workload (1-week workload) divided by chronic workload (previous 4-week average acute workload) were classified into discrete ranges by z-scores. Relative risk (RR) for each range was then calculated between injured and non-injured players using specific GPS variables: total distance, low-intensity distance, high-speed running distance, sprint distance, accelerations and decelerations.

Results The greatest non-contact injury risk was when the chronic exposure to decelerations was low (<1731) and the ACWR was >2.0 (RR=6.7). Non-contact injury risk was also 5–6 times higher for accelerations and low-intensity distance when the chronic workloads were categorised as low and the ACWR was >2.0 (RR=5.4–6.6), compared with ACWRs below this. When all chronic workloads were included, an ACWR >2.0 was associated with a significant but lesser injury risk for the same metrics, plus total distance (RR=3.7–3.9).

Conclusions We recommend that practitioners involved in planning training for performance and injury prevention monitor the ACWR, increase chronic exposure to load and avoid spikes that approach or exceed 2.0.

exposure can increase injury risk. An elevated risk of injury (relative risk [RR]=5.1) with a very high 3-weekly accumulation of accelerations (ACC) (>9254) has been demonstrated in elite youth football players.⁵ More recently, a greater absolute and relative exposure in the 3 weeks prior to injury was reported in professional football players.⁶ In contrast, other work in elite football found that gradually increasing the exposure to moderate to high training loads produced a smaller association to injury risk than exposure to lower training loads.⁷ Therefore, loads should be monitored over longer periods of time, specifically, how much is performed and how they are prescribed.

Due to the increasing physical demands of the Premier League⁸ and the congested fixture schedule at the top level, players are required to repeatedly perform high workloads. Therefore, appropriate training loads that produce adaptations to enhance their fitness levels and tolerance to physical stress are required.⁹ In this case, higher workloads would appear to be protective, while lower workloads may be insufficient to induce adaptations or result in detraining thereby increasing the risk of injury. Supporting this, Gabbett's¹⁰ training-injury prevention paradox states that excessive, rapid increases in load heighten the risk of injury, whereas chronic exposure to higher loads enhances the physical capacities of the athletes making them more resilient to injury while also enhancing performance. Consequently, there has been growing support for the acute:chronic workload ratio (ACWR) as a method of prescribing appropriate training loads. This involves the assessment of the absolute 1-week workload (acute workload) relative to 4-week chronic workload (4-week average acute workload).¹¹ A workload index can then be calculated indicating whether the individual's acute workload is greater, less than or equal to the chronic workload they have been prepared for.

Acute workload spikes have been associated with increased injury risk in football, with metres per minute prior to injury being significantly higher than the season average. However, only 16 injuries were analysed, match data were not recorded and the ACWR was not calculated, therefore warranting further research.¹² Consequently, the authors of this study investigated the relationship between the ACWR and injury risk in elite youth football players. A significantly increased risk of injury

INTRODUCTION

Typically, top-level football players sustain two injuries per season, resulting in 50 injuries within a squad of 25 players.¹ During the 2016/2017 football season, £177 million was paid out in wages to injured Premier League players, with the average wage per injury being over £248 000.² Consequently, throughout a season, clubs could be expected to pay around £12.4 million in wages alone, not including additional treatment costs, to players who are unavailable due to injury. In addition, across 24 European clubs, player availability was positively related to team success, defined by league ranking, and points per match.³

All injuries occur when an athlete is exposed to a given workload.⁴ Thus, each training or competition bout performed has the potential for athletic injury, indicating that inappropriate workload



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Bowen L, Gross AS, Gimpel M, et al. *Br J Sports Med* Epub ahead of print: [please include Day Month Year]. doi:10.1136/bjsports-2018-099422

(RR=2.6) was reported with high ACWR (1.4–1.9) for high-speed distance when the chronic workload was low (<938 m).⁵ While these findings cannot be generalised, they suggest that monitoring the ACWR in professional football may be a key injury prevention strategy.

Furthermore, most studies regarding workload–injury relationships have excluded contact injuries as they are assumed to be unavoidable. However, our previous work in youth football players found very high ACWR to be associated with contact injury risk across several workload measures (RR=4.8–5.0).⁵ It was concluded that by increasing fitness levels and limiting fatigue (ie, reducing the ACWR), players may be able to respond more quickly to avoid the rapid, unpredictable movements preceding contact injury. Therefore, the inclusion of contact injuries may provide additional insight into workload–injury relationships.

Understanding the workload–injury relationship is fundamental to optimising performance and maximising player availability. Yet, there is very limited research exploring the relationships between workloads and injury in professional football. Furthermore, despite its growing popularity as a load monitoring method, the ACWR and the associated injury risks require further exploration. Therefore, we aimed to examine the relationships of accumulated workloads, the ACWR and injury risk in Premier League football across three seasons.

METHOD

Participants

Data were collected from football players (n=33) from one English Premier League club (age: 25.4±3.1 years, stature: 182.0±6.9 cm, body mass: 79.9±7.7 kg). All players trained on a full-time basis and played competitive fixtures within the Premier League during the 2014–2015, 2015–2016 and 2016–2017 seasons. Ten (30%) participants competed in all three seasons, 8 (24%) participants competed in two seasons and the remaining 15 competed in one season, resulting in 61 individual football seasons, analysed as independent data points. Goalkeepers were excluded from the study due to the different nature of their activity.

Quantifying workload

A global positioning system (GPS) was used to quantify workload data collected from all on-pitch training sessions and friendly matches. The GPS units (Viper 2, StatSports, Ireland), were placed between the scapulae of the players in bespoke vests. These units sampled at 10 Hz and the accelerometers at 100 Hz. Following each session, the data were downloaded into the specialised analysis software (Viper, 2.1.3.0). Competitive match data were recorded using a semiautomated camera system (video (VID)) (TRACAB; ChyronHego, New York, USA) provided, as standard by the English Premier League. The raw

data files were then imported into the GPS software and analysed in an identical manner.

The validity and reliability of both GPS and VID for quantifying the physical demands of team sports has been demonstrated by numerous studies. The interchangeability of the two systems has also been established.¹³ In addition, as part of internal club research, the two systems used in this study were tested during a number of discrete tasks within a football-specific circuit, as well as during match play. Strong correlations ($r=0.7–0.9$, $p<0.05$) and small percentage differences (<10%) were found between GPS and VID for distances covered at all speeds except sprint distance, in line with previous work.¹³ Furthermore, both systems demonstrated good within player reliability (coefficient of variation (CV) <5%) and low percentage bias (<10%) when compared with the criterion during the circuit.

For sessions when data were unavailable for a participant (n=1149 of 10221; 11%) as a result of them not wearing a unit, not having match data, not completing the entire session or the data being deemed unreliable due to intermittent satellite signal, estimations were made as follows:

Main training session data: estimated by calculating squad averages for drills completed (n=607 of 10221; 6%).

International data: estimated by calculating the squad average of the other international players during the period of the international breaks¹⁴ (n=306 of 10211; 3%).

Game data: matches were only monitored using VID from 2015/2016 onwards. Prior to this, match data were estimated using individual season game averages (from a minimum of three matches) from the data collected in 2015/2016 and 2016/2017. For players where this was not available (n=6), estimations were made based on friendly matches in which GPS was worn. Game averages were extrapolated according to individual game time, as per previous work¹² (n=236 of 10221; 2%).

The variables defined in table 1 were selected for use in this study due to their relevance to running loads (and potential injury). All variables were taken from the StatSports software (Viper).

Definition of injury

Injury information was classified by the club doctor and senior chartered physiotherapists. A recordable injury was defined as one that caused any absence from future football participation, that is, a time loss injury.¹⁵ Injuries were classified as being either: minimal (1–3 days of football activity missed), mild (4–7 days of football activity missed), moderate (1–4 weeks of football activity missed) or severe (4+ weeks of football activity missed).¹⁵ Injuries were also categorised by injury type and body site. The mechanism in which a participant acquired an injury was also classified as being non-contact or contact in nature.

Table 1 Definition of workload variables

Variable	Definition
Total distance	Total covered (m): this includes walking, jogging, fast running and sprinting.
Low-intensity distance	Total distance covered (m) below 14.4 km/hour.
High-speed running distance	Total distance covered (m) between 19.8 km/hour and 25.2 km/hour.
Sprint distance	Total distance covered (m) over 25.2 km/hour.
Accelerations	An increase in GPS speed data for at least half a second with maximum acceleration in the period at least 0.5 m/s/s.
Decelerations	A decrease in GPS speed data for at least half a second with maximum deceleration in the period at least 0.5 m/s/s.
GPS, global positioning system.	

Table 2 Workload classifications and boundaries for accumulated workloads over 1–4 weeks

			No. of weeks accumulated			
	Classification	Z-score	1	2	3	4
TD (m)	Very low	≤ -2.00	11 150	24 858	37 202	45 843
	Low	-1.99 to -1.00	11 151–17 539	24 859–35 785	37 203–52 504	45 844–67 519
	Low to moderate	-0.99 to 0.00	17 540–24 041	35 786–46 733	52 505–68 677	67 520–89 707
	Moderate to high	0.00 to 0.99	24 042–30 549	46 734–57 697	68 678–84 830	89 708–111 863
	High	1.00 to 1.99	30 550–37 065	57 698–68 685	84 831–101 176	111 864–134 050
	Very high	≥ 2.00	37 066	68 686	101 177	134 051
LID (m)	Very low	≤ -2.00	9179	20 347	30 002	37 324
	Low	-1.99 to -1.00	9180–14 627	20 348–29 653	30 003–43 487	37 325–56 070
	Low to moderate	-0.99 to 0.00	14 628–20 108	29 654–39 026	43 488–57 279	56 071–74 824
	Moderate to high	0.00 to 0.99	20 109–25 644	39 027–48 423	57 280–71 110	74 825–93 845
	High	1.00 to 1.99	25 645–31 160	48 424–57 886	71 111–85 119	93 846–112 896
	Very high	≥ 2.00	31 161	57 887	85 120	112 897
HSD (m)	Very low	≤ -2.00	110	509	904	1251
	Low	-1.99 to -1.00	111–542	510–1215	905–1861	1252–2464
	Low to moderate	-0.99 to 0.00	543–979	1216–1916	1862–2827	2464–3702
	Moderate to high	0.00 to 0.99	980–1418	1917–2624	2828–3791	3703–4941
	High	1.00 to 1.99	1419–1853	2625–3326	3792–4778	4942–6176
	Very high	≥ 2.00	1854	3327	4779	6177
SD (m)	Very low	≤ -2.00				
	Low	-1.99 to -1.00	0–52	0–149	0–245	0–336
	Low to moderate	-0.99 to 0.00	53–210	150–409	246–600	337–782
	Moderate to high	0.00 to 0.99	211–368	410–672	601–955	783–1230
	High	1.00 to 1.99	369–528	673–932	956–1310	1231–1680
	Very high	≥ 2.00	529	933	1311	1681
Acc (no.)	Very low	≤ -2.00	862	1945	2832	3510
	Low	-1.99 to -1.00	863–1397	1946–2851	2833–4166	3511–5352
	Low to moderate	-0.99 to 0.00	1398–1936	2852–3753	4166–5510	5353–7193
	Moderate to high	0.00 to 0.99	1937–2472	3754–4662	5511–6855	7194–9042
	High	1.00 to 1.99	2473–3010	4663–5576	6856–8200	9043–10 902
	Very high	≥ 2.00	3011	5577	8201	10 903
Dec (no.)	Very low	≤ -2.00	794	1795	2625	3242
	Low	-1.99 to -1.00	795–1287	1796–2625	2626–3842	3243–4933
	Low to moderate	-0.99 to 0.00	1288–1782	2626–3457	3843–5073	4934–6625
	Moderate to high	0.00 to 0.99	1783–2277	3458–4292	5074–6308	6626–8323
	High	1.00 to 1.99	2278–2771	4293–5131	6309–7459	8324–10 015
	Very high	≥ 2.00	2772	5132	7460	10 016

Acc, number of accelerations; Dec, number of decelerations; HSD, high-speed distance in metres; LID, low-intensity distance in metres; SD, sprint distance in metres; TD, total distance in metres.

Data analyses

Data were categorised in weekly blocks from Monday to Sunday. Every time a player participated in a training session or match, data were analysed in two ways. First, the previous 1-weekly, 2-weekly, 3-weekly and 4-weekly loads were calculated. The loads were then classified into discrete ranges from very low through to very high using z-scores¹⁶ (table 2). The relationships between these weekly cumulative loads and subsequent injury were investigated. Injuries that occurred within the next 7 days were included for analysis.¹⁷

Second, the acute workload for the current week was calculated as the 1-week load and chronic workload as the previous 4-week rolling average acute workload. The acute and chronic workloads were uncoupled to prevent them being falsely correlated. The ACWR was calculated by dividing the acute workload by the chronic workload.¹⁸ Only acute workloads that were preceded by four complete weeks were included in the ratio calculations. A value of greater than 1 represents an acute workload greater than the chronic workload. Chronic workloads were also separated into

high and low categories by the median score for each variable.¹⁹ From this, workload–injury relationships between ACWR ratios combined with high and low chronic workloads were analysed. As with accumulated workloads, the ratios were categorised based on z-scores (table 3). Only conditions that contained 20 or more injuries were included in the statistical analysis to allow for moderate to strong associations to be made.²⁰ Consequently, data were excluded for incidences when the chronic workloads were high for both non-contact and contact injuries. This was also the case for contact injuries when the chronic loads were low.

Statistical analyses

The analysis was performed in a similar manner to the previous work of Hulin *et al.*¹⁸ Injury incidence was determined by dividing total number of injuries by the ‘on-legs’ exposure time and reported as rates per 1000 hours. Injury risks were calculated as the number of injuries sustained relative to the number of exposures to each

Table 3 Workload classifications and boundaries for: (A) acute:chronic workload ratios overall, (B) acute:chronic workload ratios combined with high chronic workloads and (C) acute:chronic workload ratios combined with low chronic workloads

	Classification	Z-score	A	B	C
TD (m)				≥22 325	<22 325
	Very low	≤−2.00	0.39	0.49	0.43
	Low	−1.99 to −1.00	0.40–0.70	0.50–0.72	0.44–0.77
	Low to moderate	−0.99 to 0.00	0.71–1.08	0.73–0.95	0.78–1.22
	Moderate to high	0.00 to 0.99	1.09–1.47	0.96–1.18	1.23–1.67
	High	1.00 to 1.99	1.48–1.86	1.19–1.42	1.68–2.13
LID (m)	Very high	≥2.00	1.87	1.43	2.14
				≥19 322	<19 322
	Very low	≤−2.00	0.35	0.43	0.40
	Low	−1.99 to −1.00	0.36–0.68	0.44–0.72	0.41–0.68
	Low to moderate	−0.99 to 0.00	0.69–1.09	0.73–1.02	0.69–1.17
	Moderate to high	0.00 to 0.99	1.10–1.50	1.03–1.32	1.18–1.66
HSD (m)	High	1.00 to 1.99	1.51–1.97	1.33–1.67	1.67–2.14
	Very high	≥2.00	1.98	1.68	2.15
				≥946	<946
	Very low	≤−2.00	0.09	0.20	0.09
	Low	−1.99 to −1.00	0.10–0.54	0.21–0.55	0.10–0.55
	Low to moderate	−0.99 to 0.00	0.55–1.08	0.56–0.93	0.56–1.17
SD (m)	Moderate to high	0.00 to 0.99	1.09–1.62	0.94–1.30	1.18–1.80
	High	1.00 to 1.99	1.63–2.16	1.31–1.68	1.81–2.47
	Very high	≥2.00	2.17	1.69	2.48
				≥174	<174
	Very low	≤−2.00	0–0.29	0–0.36	0–0.29
	Low	−1.99 to −1.00	0.30–1.16	0.37–1.01	0.30–1.35
Acc (no.)	Low to moderate	−0.99 to 0.00	1.17–2.04	1.02–1.66	1.36–2.40
	Moderate to high	0.00 to 0.99	2.05–2.92	1.67–2.39	2.41–3.46
	High	1.00 to 1.99	2.93	2.40	3.47
	Very high	≥2.00		≥1881	<1881
	Very low	≤−2.00	0.32	0.46	0.34
	Low	−1.99 to −1.00	0.33–0.67	0.47–0.71	0.35–0.73
Dec (no.)	Low to moderate	−0.99 to 0.00	0.68–1.10	0.72–0.96	0.74–1.25
	Moderate to high	0.00 to 0.99	1.11–1.54	0.97–1.21	1.26–1.77
	High	1.00 to 1.99	1.55–1.97	1.22–1.47	1.78–2.29
	Very high	≥2.00	1.98	1.48	2.30
				≥1731	<1731
	Very low	≤−2.00	0.33	0.45	0.37
	Low	−1.99 to −1.00	0.34–0.67	0.46–0.70	0.38–0.73
	Low to moderate	−0.99 to 0.00	0.68–1.10	0.71–0.95	0.74–1.26
	Moderate to high	0.00 to 0.99	1.11–1.54	0.96–1.20	1.27–1.78
	High	1.00 to 1.99	1.55–1.98	1.21–1.45	1.79–2.31
	Very high	≥2.00	1.99	1.46	2.32

Acc, number of accelerations; Dec, number of decelerations; HSD, high-speed distances in metres; LID, low-intensity distance in metres; SD, sprint distance in metres; TD, total distance in metres.

workload classification.¹⁸ Exposure data were recorded as per the procedures outlined by the Fédération de Football Association Medical Assessment Research Centre.¹⁵ A binary logistic regression model was used to compare workloads between injured and non-injured players for all workload variables independently. Accumulated workload and ACWR were independently modelled as predictor variables. RR was then calculated using to determine the magnitude of the injury risk above and below given workloads or ratios (MedCalc Software, Ostend, Belgium). When an RR was greater than 1.00, an increased risk of injury was reported (ie, RR=1.50 is indicative of a 50% increased risk) and vice versa. For an RR to be significant, 95% CIs did not contain the null RR of 1.00. Data were

analysed using IBM SPSS Statistics V.25.0 and reported as means and 95% CI. Significance was accepted at $p < 0.05$.

RESULTS

Injury incidence

For the duration of the study, 132 injuries (13.3/1000 hours 'on-legs' exposure time) were recorded (2014–2015 season, 17.6/1000 hours; 2015–2016 season, 10.2/1000 hours; 2016–2017 season, 12.4/1000 hours), including contact and non-contact injuries (see online supplementary appendix A). The knee was the most common site of injury across the three

seasons (2.9/1000 hours), 69% of which were non-contact injuries (2.0/1000 hours), predominantly meniscal or cartilage lesions and ligament sprains (0.9 and 0.7/1000 hours, respectively). The ankle was the most common site of contact injury (1.9/1000 hours), with the most common type being ligament sprains (1.6/1000 hours). The injury incidence in competition was over five times that of training (33.7/1000 hours vs 5.8/1000 hours). In particular, contact injuries were considerably greater in competition than in training (16.9/1000 hours vs 1.3/1000 hours). Despite a lower exposure to competition, 80% of contact injuries occurred in matches. The total number of days missed through injury was 4820 (36.5 ± 62.7 [mean \pm SD] days per injury).

Overall injuries

A low chronic workload of ACC (<1881), decelerations (DEC; <1731) and low-intensity distance (LID; <19222 m) combined with a very high ACWR (>2.0) elicited the greatest overall injury risk (RR=3.2, 95% CI 1.3 to 7.6, $p=0.01$; RR=3.5, 95% CI 1.5 to 8.2, $p=0.01$; and RR=2.76, 95% CI 1.2 to 6.6, $p=0.02$, respectively). The risk was also significant for very high ACWR of the same metrics, plus total distance (TD), combined with all chronic loads (RR=2.4–2.6) (table 5). Conversely, a low ACWR of TD (0.4–0.7) for all chronic loads was associated with a decreased injury risk (RR=0.2, 95% CI 0.1 to 0.8, $p=0.02$).

Non-contact injuries

Low chronic workloads combined with very high ACWRs for TD (>2.14), LID (>2.15), ACC (>2.30) and DEC (>2.32) resulted in a non-contact injury risk 5–7 times greater than ACWRs below this (RR=4.5 (TD), -6.6 (DEC), $p<0.05$). Additionally, a low amount of TD accumulated over 4 weeks (45 844–67 519 m) also resulted in an increased risk (RR=2.2, 95% CI 1.0 to 4.6, $p=0.04$) (table 4). Significant risks were also found for TD, ACC, DEC and LID for all chronic loads when the ACWR was very high (RR=3.7–3.9) (table 5).

Contact injuries

Moderate to high ACWR (1.1–1.5) for TD, DEC and LID produced the largest contact injury risk (RR=2.0, 95% CI 1.0 to 4.0, $p=0.04$; RR=2.0, 95% CI 1.0 to 4.0, $p=0.04$; and RR=2.6, 95% CI 1.3 to 5.2, $p=0.01$, respectively). A moderate to high amount of TD (24 042–30 549 m) and a low to moderate amount of DEC (1288–1782) accumulated over a week also showed a heightened risk of contact injury (RR=2.1, 95% CI 1.1 to 4.0, $p=0.03$ and RR=2.0, 95% CI 1.1 to 3.9, $p=0.03$, respectively).

DISCUSSION

This is the first study to explore the relationship of both accumulated GPS-derived loads and ACWR with contact and non-contact injury risk at an English Premier League football club. This extends our previous work,⁵ and that of others,^{7 18 19 21 22} showing that a number of GPS-derived workloads were associated with injury risk.

Non-contact injuries and workload

A very high ACWR combined with low chronic workload categories demonstrated the greatest non-contact injury risk for most metrics (except high-speed distance [HSD] and sprint distance [SD]), with DEC being most strongly associated with RR (ACWR >2.3 , RR=6.6). When all chronic workloads were analysed, a very high ACWR demonstrated a lesser but still significant risk

for the same metrics (RR=3.7–3.9). These findings are in line with studies in cricket,¹⁸ rugby,^{19 22} Australian football,²³ Gaelic football²⁴ and football⁵ where high ACWR, referred to as 'spikes' in workload, have been associated with heightened injury risk. Due to the inevitable increased exposure to risk with greater workloads, previous research has focused on the higher load–higher injury risk relationship.^{7 25} However, the above findings, alongside the lack of significant risks associated with high accumulated loads in this study, support a growing body of literature suggesting that acute, excessive, rapid increases in loads may be responsible for a large proportion of non-contact injuries, rather than chronic exposure to higher loads.¹⁰

The protective effect of high chronic loads versus low chronic loads has been reported in rugby.^{19 22} They concluded that the players who were capable of achieving high exposure had the enhanced physical attributes needed for decreased injury risk. This theory has recently been demonstrated in hurling, where players with well-developed lower body strength, repeated sprint ability and speed tolerated higher training loads and had a reduced risk of injury compared with lower performance groups.²⁶ In youth football, a high ACWR combined with low chronic HSD load showed a significantly increased risk of non-contact injury (ACWR=1.4–2.0, RR=2.6), which was not evident when combined with high chronic HSD (ACWR=1.3–1.8, RR=0.5).⁵

In the current study, there were not enough injuries when the chronic workloads were high to determine the RRs of non-contact injury, further indicating a potential protective effect. Ultimately, training at higher workloads may cause players to develop a greater tolerance for the increasing intensity and fatigue of competition.¹⁰ Concurrently, reducing training loads, while lowering a player's exposure to risk, may also have a negative effect on fitness and physical preparedness, potentially increasing the risk. Therefore, as per the training–injury prevention paradox model,¹⁰ optimal load management to minimise injury risk should involve appropriate, progressive exposure to higher loads while avoiding training load spikes that the player is not prepared for.

Contact injuries and workload

Very little research has investigated the relationship between contact injuries and workload, despite early workload–injury research suggesting that players with better developed physical capacity may be at less risk of contact injury.²⁷ In the current study, 80% of all contact injuries occurred in matches, similar to previous injury incidence reports.²⁸ This may be due to the high speed and intensity of play, resulting in more body contact such as sliding and tackling.²⁹ The risk of contact injury was greatest when the ACWR was moderate to high for TD, LID and DEC (RR range=2.0–2.6), meaning the acute workload was very similar to the chronic workload.

Also, as load was categorised by z-scores, contact risk was highest for the most commonly occurring ratios ($z=0.0$ – 1.0). Players who are regularly in the team, and therefore more at risk of contact injury, typically have a lack of variation in their workload due to a large proportion of the weekly load being attained from matches. Therefore, when the match load was constant, variations in the training load produced very little fluctuation in the total acute load.²³ Thus, it would appear that in the current study, contact injury is most likely to be related to match exposure, rather than the prescribed workload. The lack of association of the ACWR to contact injuries is highlighted further by the

Table 4 Injury risk associated with accumulated workloads over 1–4 weeks

No. of weeks accumulated		1			2			3			4		
		NC	C	Overall	NC	C	Overall	NC	C	Overall	NC	C	Overall
TD (m)	Very low	0.49	0.67	0.28	1.38	0.94	1.19	1.55	2.17	1.81	0.92	1.28	1.07
	Low	0.94	0.16	0.59	1.11	1.01	1.07	1.26	0.32	0.84	2.18*	0.66	1.49
	Low to moderate	1.51	0.85	1.20	1.03	0.86	0.95	1.00	0.69	0.86	0.97	0.63	0.82
	Moderate to high	0.80	2.09*	1.22	1.10	1.73	1.33	0.96	1.55	1.18	0.81	1.79	1.13
	High	0.84	0.93	0.88	0.57	0.18	0.40*	0.66	0.68	0.67	0.92	0.48	0.73
	Very high	1.01	1.39	1.17	1.29	1.78	1.50	1.52	2.11	1.77	0.86	2.57	1.07
LID (m)	Very low	0.40	0.55	0.23	1.32	0.89	1.14	0.51	2.23	1.20	0.91	1.26	1.06
	Low	0.81	0.34	0.60	1.15	1.34	1.23	1.55	0.32	1.00	1.27	0.66	1.00
	Low to moderate	1.17	1.11	1.15	0.98	1.19	1.06	1.06	0.88	0.98	1.36	0.95	1.18
	Moderate to high	1.14	1.65	1.34	1.15	1.13	1.14	0.77	1.30	0.97	0.70	1.32	0.93
	High	0.84	0.93	0.88	0.58	0.38	0.49	1.02	0.92	0.98	1.07	0.70	0.91
	Very high	1.01	0.65	0.58	1.18	0.76	0.68	2.63	0.83	1.50	0.86	1.18	0.50
HSD (m)	Very low	2.61	3.57	1.52	3.23	1.01	1.83	0.44	0.60	0.25	1.43	0.97	1.23
	Low	1.15	0.17	0.70	1.46	0.59	1.07	1.52	0.85	1.22	1.49	0.70	1.14
	Low to moderate	0.86	1.07	0.94	0.92	1.58	1.16	0.99	1.36	1.34	0.87	1.00	0.92
	Moderate to high	1.16	1.33	1.23	0.89	0.99	0.93	0.74	0.79	0.76	0.98	0.82	0.91
	High	1.33	1.61	1.44	0.76	0.40	0.60	1.59	0.89	1.28	0.65	1.46	1.19
	Very high	0.24	0.68	0.28	0.78	2.22	1.37	0.36	2.18	0.89	0.49	2.45	1.00
SD (m)	Very low												
	Low	1.10	0.13	0.60	1.20	0.78	1.02	1.29	0.44	0.88	1.83	0.65	1.29
	Low to moderate	1.16	1.31	1.22	1.16	0.90	1.04	1.00	1.19	1.04	1.06	1.22	1.13
	Moderate to high	1.08	1.16	1.12	1.01	1.24	1.10	1.23	0.72	0.98	0.98	0.59	0.62
	High	0.82	0.85	0.83	0.41	0.87	0.60	0.36	1.76	1.16	0.65	1.39	1.14
	Very high	0.23	1.36	0.55	0.96	1.34	1.12	0.97	1.32	0.82	0.49	2.15	1.16
Acc (no.)	Very low	0.45	0.61	0.26	1.18	0.80	1.02	0.48	2.10	1.13	1.28	1.16	1.23
	Low	0.93	0.33	0.67	1.18	1.06	1.13	1.50	0.64	1.12	1.06	0.71	0.91
	Low to moderate	1.08	1.62	1.29	1.31	1.89	1.53*	1.10	0.92	1.02	0.99	1.00	0.99
	Moderate to high	1.28	0.74	1.03	0.85	0.68	0.78	0.81	1.10	0.92	1.04	1.41	1.18
	High	0.70	1.45	0.99	0.56	0.37	0.48	1.02	0.67	0.87	0.85	0.22	0.57
	Very high	1.12	1.54	1.30	1.71	2.36	1.98	2.14	2.96	2.48	1.39	4.24	1.76
Dec (no.)	Very low	0.50	0.69	0.29	1.32	0.89	1.14	0.48	0.72	1.15	1.39	1.17	0.98
	Low	1.04	0.32	0.71	0.91	1.61	1.19	1.40	0.60	1.05	1.09	0.73	0.94
	Low to moderate	0.89	2.04*	1.29	1.11	1.34	1.20	1.21	0.92	1.08	1.03	1.08	1.05
	Moderate to high	1.31	0.55	0.94	1.13	0.79	0.97	0.69	1.14	0.86	1.08	1.31	1.17
	High	0.84	1.46	1.09	0.55	0.37	0.47	1.18	0.65	0.95	0.84	0.22	0.56
	Very high	1.06	1.47	1.23	1.64	2.27	1.91	2.04	2.82	2.37	1.30	3.96	1.64

Acc, number of accelerations; Dec, number of decelerations; HSD, high-speed distances in metres; LID, low-intensity distance in metres (m); SD, sprint distance in metres; TD, total distance in metres; N, non-contact injury; C, contact injury, *, $p < 0.05$; **, $p < 0.01$.

large RR of non-contact injury following an acute spike compared with overall injury, suggesting that including contact injuries reduces the association of the ACWR with injury risk. Consequently, these injuries should be analysed separately when establishing workload-injury relationships and determining uniform injury definitions across research.³⁰

Potential limitations

Previous studies have highlighted the limitations of using estimated match data, as it does not account for match to match variability.¹⁴ This study has attempted to improve on this by including match data for the 2015/2016 and 2016/2017 seasons. However, for the 2014/2015 season, TRACAB data were not available, resulting in estimations being calculated as

Table 5 Injury risks associated with (A) acute:chronic workload ratios overall, (B) acute:chronic workload ratios combined with high chronic workloads and (C) acute:chronic workload ratios combined with low chronic workloads

		A			B			C					
		NC	C	Overall	NC	C	Overall	NC	C	Overall			
TD (m)	Very low							2.04					
	Low	0.35	0.11		0.19*			0.16	0.59			0.35	
	High	0.84	1.46	1.09	0.55	0.37	0.47	1.18	0.65	0.95	0.84	0.22	0.56
	Very high	1.06	1.47	1.23	1.64	2.27	1.91	2.04	2.82	2.37	1.30	3.96	1.64
	Low to moderate	1.32	0.69	1.01				1.01		1.56		1.22	
	Moderate to high	0.91	2.03*	1.31				1.18		0.48		0.90	
	High	0.29	1.72	0.87				1.86		0.50		0.96	
	Very high	3.67*	0.88	2.40*				0.80		4.50*		2.61	
LID (m)	Very low							2.79					
	Low	0.39	0.25	0.33				0.30		0.25		0.16	
	Low to moderate	0.83	0.48	0.66				0.80		1.15		1.00	
	Moderate to high	1.52	2.60*	1.91*				2.08*		0.85		1.13	
	High	0.16	1.40	0.57				0.59		0.23		0.58	
	Very high	3.93*	0.94	2.56*						5.39*		2.76*	
HSD (m)	Very low												
	Low	1.35	0.26	0.85				0.80		1.67		0.97	
	Low to moderate	0.92	1.64	1.18				0.84		1.73		1.78	
	Moderate to high	1.20	0.80	1.02				1.36		0.70		0.72	
	High	0.52	1.07	0.75				1.36		0.39		0.48	
	Very high	0.66	0.88	0.76				0.38		0.39		0.51	
SD (m)	Very low												
	Low	1.05	0.15	0.57				0.24		1.25		0.75	
	Low to moderate	0.72	1.40	0.96				1.68		0.66		0.83	
	Moderate to high	1.10	1.12	1.11				0.65		1.31		1.40	
	High	1.36	0.86	1.14				1.78		1.34		1.10	
	Very high	1.96	0.82	1.46				0.58		1.01		0.63	
Acc (no.)	Very low							3.50					
	Low	0.42	0.27	0.35				0.35		0.18		0.11	
	Low to moderate	0.79	0.80	0.79				0.84		1.54		1.49	
	Moderate to high	1.40	1.81	1.57*				1.37		0.60		0.75	
	High	0.33	0.90	0.57				1.27		0.28		0.71	
	Very high	3.86*	0.92	2.52*				1.12		5.90**		3.18*	
Dec (no.)	Very low							3.59					
	Low	0.44	0.28	0.37				0.18		0.16		0.09	
	Low to moderate	0.88	0.80	0.85				0.98		1.41		1.40	
	Moderate to high	1.23	1.99*	1.52				1.31		0.71		0.86	
	High	0.34	0.45	0.39				1.29		0.25		0.64	
	Very high	3.73*	0.89	2.44*				0.88		6.58**		3.47*	

Acc, number of accelerations; C, contact injury; Dec, number of decelerations; HSD, high-speed distances in metres; LID, low-intensity distance in metres; N, non-contact injury; SD, sprint distance in metres; TD, total distance in metres; *, $p < 0.05$; **, $p < 0.01$.

per previous work,¹² emulating the aforementioned limitation. Additionally, as the match data for the latter two seasons were collected using a different system than training, the precision and sensitivity of the data may be decreased, despite it being calibrated to maximise between system agreements.¹³ With technological advancements, and the recent admittance of GPS in league matches, future research should aim to use a single monitoring system for both competition and training.

One potential explanation for the lack of significant non-contact injury risk for very high acute HSD and SD, despite all other metrics reporting otherwise, may be the use of absolute speed thresholds in this study. Buchheit¹⁴ recently stated that the use of fixed thresholds may reduce the sensitivity of the ACWR due to the varying locomotor profiles between players, particularly as subtle differences in speed at high intensity have been found

to have important implications on injury risk. Future research could consider the use of individualised thresholds, although caution must be taken when anchoring all locomotor categories to one fitness measure.³¹

The present study only examined external load; however, the incorporation of the rating of perceived exertion (RPE) values, as a measure of internal workload, may provide a more complete insight into the likelihood of injury, as well as taking into consideration the athlete's response to a given workload.¹⁷ Fanchini *et al*²¹ recently analysed the ACWR in relation to injury risk in elite football using RPE as their load measure. Similar to the current study and the previous work of the authors,⁵ they demonstrated a heightened risk with acute 'spikes'. A combination of both methods may give a more complete assessment of load-related risk, while also considering

the validity and specificity of the chosen metrics to the sport and the individual.³²

Calculating the ACWR using rolling averages is evidence based and supported by a large body of literature. However, future research may consider using exponentially weighted moving averages, which consider the decaying nature of fitness and fatigue over time. This method has recently been shown to have a greater sensitivity to increases in injury risk at higher ACWRs.³³

The statistical power of this study was not calculated prospectively. As retrospective power analysis calculations are not appropriate,³⁴ the power analysis was not included. However, this study included 81 injury cases, which is enough to make moderate to strong associations regarding injury risk factors.²⁰ Future studies should ensure prospective power analysis for inclusion. Furthermore, as commonly recommended in elite sport research, future work involving multiple clubs would enhance the ability to generalise these findings, advance the statistical analysis and detect small to moderate associations (+200 injury cases).

SUMMARY AND CONCLUSIONS

In summary, ACWR had a stronger association to non-contact injury risk in this cohort of English Premier League football players than accumulated loads, suggesting the rapid increase in load is more indicative of injury than the cumulative amount of load performed. Specifically, very high acute spikes when the chronic loads were low corresponded to the greatest non-contact injury risk. We recommend that training programmes should involve progressive exposure to higher loads to enhance physical capacities while minimising the risks associated with rapid, excessive spikes. Due to the majority of contact injuries occurring during competition, which is both inevitable and relatively non-modifiable by practitioners, it is unlikely that they were associated with a given workload. While this study provides an initial insight into the relationships between workload and injury risk, care should be taken when applying the findings beyond the studied population.

Data were excluded for categories where there were less than 20 injuries, as moderate to strong associations between workload and injury could not be made.²⁰

What are the findings?

- For English Premier League football players, when chronic load exposure is low, an acute:chronic workload ratio (ACWR) >2 is associated with 5–7 times the risk of a soft tissue injury than players whose ACWR is below this.
- Regardless of chronic exposure, a spike in workload is associated with increased injury risk, although this risk is lessened with greater chronic loads.
- A spike in decelerations was associated with the greatest non-contact injury risk.

How might it impact on clinical practice in the future?

- Our data provide initial recommendations for monitoring ACWRs in English Premier League football players.
- We highlight some global positioning system-derived variables that may serve practitioners who monitor elite football players' workload to prevent injury.

Contributors All authors listed met the conditions required for full authorship. LB designed the initial study proposal, which was presented to ASG, SB-L and MG. From there, all five authors met regularly to ensure a scientifically sound study design was created. LB collated and analysed the data with the help of both ASG and MG throughout the process. F-XL and SB-L advised and checked the statistical analysis. LB wrote the main body of the article, which was revised multiple times by all authors before being approved for publication. LB modified the article following revisions with contributions from SB-L. All authors are fully aware and understanding of the findings of the study and confident in the integrity of the research.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Approval was obtained from the University of Birmingham Ethical Review Committee.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement All data analysed are included in the article.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- 1 Ekstrand J, Häggglund M, Waldén M. Injury incidence and injury patterns in professional football: the UEFA injury study. *Br J Sports Med* 2011;45:553–8.
- 2 <https://www.jltsspecialty.com/media-centre/press-release/2017/june/injury-costs-for-premier-league-clubs> (accessed 20 Mar 2018).
- 3 Häggglund M, Waldén M, Magnusson H, et al. Injuries affect team performance negatively in professional football: an 11-year follow-up of the UEFA Champions League injury study. *Br J Sports Med* 2013;47:738–42.
- 4 Windt J, Gabbett TJ. How do training and competition workloads relate to injury? The workload-injury aetiology model. *Br J Sports Med* 2017;51:428–35.
- 5 Bowen L, Gross AS, Gimpel M, et al. Accumulated workloads and the acute:chronic workload ratio relate to injury risk in elite youth football players. *Br J Sports Med* 2017;51:452–9.
- 6 Lu D, Howle K, Waterson A, et al. Workload profiles prior to injury in professional soccer players. *Sci Med Football* 2017;1:237–43.
- 7 Malone S, Owen A, Newton M, et al. The acute:chronic workload ratio in relation to injury risk in professional soccer. *J Sci Med Sport* 2017;20:561–5.
- 8 Barnes C, Archer DT, Hogg B, et al. The evolution of physical and technical performance parameters in the English Premier League. *Int J Sports Med* 2014;35:1095–100.
- 9 SELYE H. The general adaptation syndrome and the diseases of adaptation. *J Clin Endocrinol Metab* 1946;6:117–230.
- 10 Gabbett TJ. The training-injury prevention paradox: should athletes be training smarter and harder? *Br J Sports Med* 2016;50:273–80.
- 11 Banister E, Calvert T, Savage M, et al. A systems model of training for athletic performance. *Aus J Sports Med* 1975;7:57–61.
- 12 Ehrmann FE, Duncan CS, Sindhusake D, et al. GPS and injury prevention in professional soccer. *J Strength Cond Res* 2016;30:360–7.
- 13 Buchheit M, Allen A, Poon TK, et al. Integrating different tracking systems in football: multiple camera semi-automatic system, local position measurement and GPS technologies. *J Sports Sci* 2014;32:1844–57.
- 14 Buchheit M. Applying the acute:chronic workload ratio in elite football: worth the effort? *Br J Sports Med* 2017;51:1325–7.
- 15 Fuller C, Ekstrand J, Junge A, et al. Consensus statement on injury definitions and data collection procedures in studies of football (soccer) injuries. *Br J Sports Med* 2006;16:83–92.
- 16 Wang Y, Chen HJ. Use of percentiles and z-scores in anthropometry. In: Preedy VR, ed. *Handbook of anthropometry: physical measures of human form in health and disease*. New York, NY: Springer Science and Business Media, 2012:29–48.
- 17 Piggott B, Newton M, McGuigan M. The relationship between training load and incidence of injury and illness over a pre-season at an Australian football league club. *J Aust Strength Cond Res* 2008;17:4–17.
- 18 Hulin BT, Gabbett TJ, Blanch P, et al. Spikes in acute workload are associated with increased injury risk in elite cricket fast bowlers. *Br J Sports Med* 2014;48:708–12.
- 19 Hulin BT, Gabbett TJ, Lawson DW, et al. The acute:chronic workload ratio predicts injury: high chronic workload may decrease injury risk in elite rugby league players. *Br J Sports Med* 2016;50:231–6.
- 20 Bahr R, Holme I. Risk factors for sports injuries—a methodological approach. *Br J Sports Med* 2003;37:384–92.

- 21 Fanchini M, Rampinini E, Riggio M, *et al.* Despite association, the acute:chronic work load ratio does not predict non-contact injury in elite footballers. *Sci Med Football* 2018;2:108–14.
- 22 Hulin BT, Gabbett TJ, Caputi P, *et al.* Low chronic workload and the acute:chronic workload ratio are more predictive of injury than between-match recovery time: a two-season prospective cohort study in elite rugby league players. *Br J Sports Med* 2016;50:1008–12.
- 23 Stares J, Dawson B, Peeling P, *et al.* Identifying high risk loading conditions for in-season injury in elite Australian football players. *J Sci Med Sport* 2018;21:46–51.
- 24 Malone S, Roe M, Doran DA, *et al.* Aerobic fitness and playing experience protect against spikes in workload: the role of the acute: chronic workload ratio on injury risk in elite Gaelic football. *Int J Sports Physiol Perform* 2016;24:1–25.
- 25 Gabbett TJ, Jenkins DG. Relationship between training load and injury in professional rugby league players. *J Sci Med Sport* 2011;14:204–9.
- 26 Malone S, Hughes B, Doran DA, *et al.* Can the workload–injury relationship be moderated by improved strength, speed and repeated-sprint qualities? *J Sci Med Sport* 2019;22:29–34.
- 27 Gabbett TJ. The development and application of an injury prediction model for noncontact, soft-tissue injuries in elite collision sport athletes. *J Strength Cond Res* 2010;24:2593–603.
- 28 Hawkins RD, Fuller CW. A prospective epidemiological study of injuries in four English professional football clubs. *Br J Sports Med* 1999;33:196–203.
- 29 Wong P, Hong Y. Soccer injury in the lower extremities. *Br J Sports Med* 2005;39:473–82.
- 30 Hulin BT. The never-ending search for the perfect acute:chronic workload ratio: what role injury definition? *Br J Sports Med* 2017;51:991–2.
- 31 Hunter F, Bray J, Towilson C, *et al.* Individualisation of time-motion analysis: a method comparison and case report series. *Int J Sports Med* 2015;36:41–8.
- 32 Blanch P, Gabbett TJ. Has the athlete trained enough to return to play safely? The acute:chronic workload ratio permits clinicians to quantify a player's risk of subsequent injury. *Br J Sports Med* 2016;50:471–5.
- 33 Murray NB, Gabbett TJ, Townshend AD, *et al.* Calculating acute:chronic workload ratios using exponentially weighted moving averages provides a more sensitive indicator of injury likelihood than rolling averages. *Br J Sports Med* 2017;51:749–54.
- 34 Zumbo BD, Hubley AM. A note on misconceptions concerning prospective and retrospective power. *J R Stat Soc Ser D Stat* 1998;47:385–8.